

CH 7.2: TRIG INTEGRALS

** We will be asked to compute some more involved **TRIGONOMETRIC INTEGRALS**. We will learn a technique of using **TRIG IDENTITIES** to rewrite the integrand, which will then enable us to compute the integral. First let's review some important facts about Trig functions.

PART 1. SOME TRIG IDENTITIES

PYTHAGOREAN IDENTITIES.

$$\begin{aligned}\cos^2\theta + \sin^2\theta &= 1 \\ \tan^2\theta + 1 &= \sec^2\theta \\ 1 + \cot^2\theta &= \csc^2\theta\end{aligned}$$

RECIPROCAL IDENTITIES

$$\begin{aligned}\csc(\theta) &= \frac{1}{\sin(\theta)} \\ \sec(\theta) &= \frac{1}{\cos(\theta)} \\ \cot(\theta) &= \frac{1}{\tan(\theta)}\end{aligned}$$

DOUBLE ANGLE IDENTITIES

$$\begin{aligned}\sin(2\theta) &= 2\sin\theta\cos\theta \\ \cos(2\theta) &= 2\cos^2\theta - 1 \\ &= \cos^2\theta - \sin^2\theta\end{aligned}$$

PRODUCT TO SUM FORMULAS

$$\begin{aligned}\sin(A)\cos(B) &= \frac{1}{2}[\sin(A-B) + \sin(A+B)] \\ \sin(A)\sin(B) &= \frac{1}{2}[\cos(A-B) - \cos(A+B)] \\ \cos(A)\cos(B) &= \frac{1}{2}[\cos(A-B) + \cos(A+B)]\end{aligned}$$


HALF-ANGLE IDENTITIES

$$\begin{aligned}\cos^2\theta &= \frac{1 + \cos(2\theta)}{2} \\ \sin^2\theta &= \frac{1 - \cos(2\theta)}{2}\end{aligned}$$

PART 2: THE UNIT CIRCLE

To evaluate $\sin(\theta)$ and $\cos(\theta)$ at a particular value of θ , the unit circle comes in handy!

- Find the correct angle θ on the unit circle.
- The point will have an ordered pair associated with it.
- The **X VALUE** represents $\cos(\theta)$
- The **Y VALUE** represents $\sin(\theta)$



PART 3: TRIG INTEGRALS of Form $\int \cos^n(x) dx$, $\int \sin^n(x) dx$

or $\int \sin^m(x) \cos^n(x) dx$

Ex 1. Evaluate the following definite and indefinite integrals:

A $\int \cos^2 \theta d\theta$

! EVEN POWER ON COSINE (NO ODD POWER)

sol:

! If this were "definite" use fundamental of calculus.

$$\int_0^{\pi/2} \cos^2 \theta d\theta$$

B $\int \cos^3 \theta d\theta$

! ODD POWER ON COSINE

sol:

*
C $\int \sin^4(t) dt$

sol:

⚠ EVEN POWER ON SINE (NO ODD POWER)

D $\int \sin(x) \cos^3(x) dx$

sol:

E $\int 2(1 + \sin \theta)^2 d\theta$

sol:

E $\int \frac{\cos^3(\sqrt{x})}{\sqrt{x}} dx$

sol:

F $\int \sin^5(x) \cos^2(x) dx$

sol:

 ODD POWER ON SINE

Ex 2. Set up an integral to express the area between the graphs
 $f(x) = \sin^3(x)$ on the interval $[0, \pi]$

and

sol:

MORE EXAMPLES

① $\int \tan(x) \sec^4(x) dx$

sol:

② $\int \tan(x) \cdot \sec^3(x) dx$

sol:

RECALL $\frac{d}{dx} [\sec(x)] = \sec(x) \tan(x)$